

Contents in full

The Authors	v
A Note from the Publisher on Primers in Biology	vii
Preface	ix
Acknowledgements	xii
CHAPTER 1 Overview of the Immune Response	
1-0 Overview: The Protective Mechanisms of Immunity	2
The immune system protects us against infectious organisms	
Immune mechanisms are divided into those of innate immunity and those of adaptive immunity	
An antigen is operationally defined	
1-1 Cells of the Immune System: Differentiation in the Bone Marrow	4
Cytokines produced by bone marrow stromal cells drive the differentiation of immune cells from their common hematopoietic precursor	
Reserves of hematopoietic cells are rapidly mobilized from the bone marrow during stress	
1-2 Cells of the Immune System: Functional Characteristics	6
Immune cells may differentiate in several stages	
Neutrophils, macrophages and dendritic cells are phagocytic cells	
Basophils, mast cells and eosinophils release inflammatory and cytotoxic mediators from intracellular stores	
Lymphocytes detect antigen by means of variable receptors	
1-3 Macrophage and Dendritic Cell Subsets	8
Macrophages and dendritic cells are versatile cells of innate immunity	
Macrophages have distinct specializations in different tissues	
Dendritic cells can vary widely in form and function	
1-4 Clonal Selection of Antigen-Specific Lymphocytes	10
The versatility and selectivity of adaptive immunity are guaranteed by selective processes acting on a highly variable receptor repertoire	
Variable receptors for antigen are generated by rearrangement of lymphocyte DNA during ontogeny	
Selection by antigen leads to self tolerance and to the specific recognition of almost any foreign organism or particle	
1-5 Major Histocompatibility Molecules and the Detection of Infection	12
T lymphocytes recognize fragments of antigen carried to the cell surface by MHC molecules	
MHC molecules are surface indicators of the proteins present in the interior compartments of cells	
Cell surface markers of different T cell classes reflect the differential recognition of MHC class I and class II molecules	
1-6 The Lymphoid System and Lymphocyte Circulation	14
There are two types of lymphoid tissues	
Secondary lymphoid tissue brings antigen together with lymphocytes	
Chemokines direct the migration of lymphocytes and determine the specialized microenvironments in which they are activated	
Secondary lymphoid tissue is sustained and can be induced by signals from immune cells	
1-7 Architecture of Secondary Lymphoid Tissues	16
Dendritic cells and follicular dendritic cells collect antigen for recognition by T cells and B cells	
The spleen is a critical filter for antigen in blood	
Antigen enters mucosal lymphoid tissues through specialized cells	
1-8 The Course of an Immune Response	18
Innate immunity and specialized classes of lymphocytes provide front-line defense against invading microorganisms	

T lymphocytes activated by dendritic cells differentiate and migrate to B cell follicles and sites of infection

Activated B lymphocytes differentiate into plasma cells and secrete antibody

At the end of an immune response, the vast majority of the antigen-specific cells die and surviving cells become resting memory cells

CHAPTER 2 Signaling and Adhesive Molecules of the Immune System

2-0 Overview: Immune Cell Surface and Signaling Molecules	22
Immune cell migration and communication are controlled by specialized cell surface and signaling molecules	
Adhesion molecules have important roles in promoting cell migration and cell–cell interaction	
Large families of signaling molecules control nearly all aspects of immune system development and function	
2-1 The Immunoglobulin Superfamily: Structural Features	24
The Ig-like domain is an important element of structure for many proteins of the immune system	
The Ig superfamily contains a diverse group of molecular recognition elements of the immune system	
2-2 Signaling by Immunoglobulin Superfamily Receptors	26
Ig superfamily members mediate antigen recognition and many functional effects of antibodies	
Activating Ig superfamily receptors signal via intracellular tyrosine kinases	
Immune cells express Ig superfamily inhibitory receptors that restrain immune responses	
2-3 Adhesion Molecules of the Immunoglobulin Superfamily	28
Ig superfamily members have important roles in cell–cell adhesion of immune cells	
The CD2 subfamily participates in the adhesion and activation of immune system cells	
Several Ig superfamily members serve as cell-bound ligands for integrins	
2-4 Integrins in Immune Function	30
Integrins mediate cell–cell and cell–matrix contacts	
The adhesive properties of integrins are modulated by other signals coming into the cell	
Integrin engagement regulates cell shape and also sends signals that support cell activation and survival	
Integrins are differentially expressed to support specializations in immune cell function	
2-5 C-Type Lectins and Carbohydrate Recognition	32
Carbohydrate-recognizing molecules have diverse roles in the immune system	
2-6 Cytokines and Cytokine Receptors	34
Polypeptide mediators are a principal means of regulating immune responses	
Cytokines act on many cell types in overlapping and coordinated ways	
Cytokines can be grouped into structurally related families that act via homologous receptors	
2-7 Cytokine Receptors that Signal via the Jak–STAT Pathway	36
The family of type I cytokines is recognized by a conserved family of receptors	
Interferons and IL-10 are related to type I cytokines and are also recognized by receptors signaling via the Jak–STAT pathway	
Type I and type II cytokine receptors signal principally via the Jak–STAT signaling pathway	
STATs are responsible for much of the specificity of cytokine receptor signaling	
Jak–STAT signaling is negatively regulated by SOCS proteins	
2-8 Shared Subunits in Subfamilies of Type I Cytokine Receptors	38
Many cytokines and hematopoietins are recognized by structurally related receptors characterized by shared receptor subunits	

2-9	The TNF Superfamily of Cell Regulators	40
	Members of the TNF superfamily regulate a variety of immune and developmental events	
	TNF-receptor superfamily members signal on oligomerization	
2-10	NF-κB and Inflammatory Cytokine Action	42
	NF- κ B activates inflammatory gene expression	
	NF- κ B is regulated by inhibitory subunits	
	Some NF- κ B heterodimers are self-inactivated	
2-11	Molecular Control of Apoptosis	44
	Apoptosis has many important roles in immune function	
	Apoptosis is triggered by a proteolytic cascade involving caspases	
2-12	Regulation of Apoptosis by Bcl-2 Family Members	46
	Increased mitochondrial outer-membrane permeability is the major trigger of the intrinsic apoptosis pathway	
2-13	Chemokines and Chemokine Receptors	48
	Chemokines direct the migration of immune-system cells	
2-14	Chemokines, Integrins and Selectins and Immune-Cell Homing	50
	Chemokines collaborate with other specialized molecules in directing the homing of lymphocytes and leukocytes to tissues	
	Selectins are C-type lectins that bind to carbohydrate structures usually on cell surfaces	
	Circulating lymphocytes express a regulatable receptor that directs their exit from lymphoid tissues	
 CHAPTER 3 Innate Immunity		
3-0	Overview: Evolution and Function of Innate Immunity	54
	Multicellular organisms have conserved mechanisms of innate immune recognition	
	Innate immunity remains effective despite rapid evolution of microbes and viruses	
	Innate immune recognition mechanisms predate adaptive immunity and instruct it	
3-1	Barriers to Infection at the Epithelium	56
	The epithelial cell layer provides the first barrier to most infections	
	Antimicrobial peptides are a widespread defense mechanism	
3-2	Innate Recognition by Soluble Collectins and Ficolins	58
	Collectins and ficolins activate innate immune protective mechanisms at mucosal surfaces and in the bloodstream	
	Collectins and ficolins have a common architecture but distinct recognition domains	
	Two lung surfactants are mucosal collectins	
	Mannose-binding lectin provides innate immune protection against some bacterial infections	
3-3	Overview of the Complement System	60
	The complement cascade links soluble factor recognition of microbes to mechanisms for their destruction	
	Inherited deficiencies in complement components lead to severe bacterial infections or to susceptibility to immune complex disease	
3-4	Activation of the Complement Cascade	62
	Initiation of complement activation by the lectin and classical pathways has substantially common features	
	Complement activation is largely confined to cell surfaces	
	The alternative pathway of complement activation provides an amplification loop that is activated at cell surfaces	
3-5	Complement Effector Actions and Receptors	64
	The complement cascade promotes inflammation, phagocytosis, membrane attack and antibody production	
	Complement receptors mediate many of the actions of complement	

3-6 Regulation of Complement and Evasion by Microbes	66
Complement activation is regulated by soluble factors and membrane-bound inhibitors	
Many bacteria and viruses have evolved mechanisms for evading complement action	
3-7 Recognition of Microbes by Phagocytes	68
Phagocytosis is a major mechanism for the destruction of microbes	
Phagocytes have receptors for recognizing and internalizing microbes	
3-8 Mechanisms of Phagocytosis	70
Phagocytosis proceeds by three distinct pathways	
Phagocytosis of apoptotic cells is usually antiinflammatory	
Inhibitory receptors can regulate phagocytosis	
3-9 Destructive Mechanisms of Phagocytes	72
Phagocytes have many mechanisms for killing internalized microbes	
Reactive oxygen and nitrogen species are used to kill internalized microorganisms	
3-10 The Toll-Like Receptor Family of Innate Immune Receptors	74
Recognition of conserved microbial components by Toll-like receptors leads to inflammation and activation of sentinel immune cells	
Accessory molecules aid Toll-like receptor recognition of some ligands	
Mammalian Toll-like receptors recognize their ligands on the cell surface or intracellularly	
3-11 Toll-Like Receptor Signaling to Cytokine Production	76
Mammalian Toll-like receptors activate the production of proinflammatory cytokines and interferon	
Mammalian Toll-like receptors signal via two major pathways	
3-12 Intracellular Sensors of Bacterial Peptidoglycan Components	78
NOD1, NOD2 and cryopyrin are intracellular sensors of peptidoglycan substructures	
The NOD domain mediates ligand-induced signal generation	
3-13 Inflammation: Initiation of an Inflammatory Response	80
Innate immune recognition triggers an inflammatory response that focuses the immune system on the site of infection	
Cytokines and lipid mediators are important inducers of inflammation	
3-14 Inflammation: Recruitment of Immune Cells	82
Leukocyte extravasation is a multi-step process	
Chemokines and small-molecule chemoattractants direct phagocytes to the site of infection	
3-15 Inflammation: Regulation and Systemic Effects	84
Inflammatory responses can be divided into two phases: the acute phase and the chronic phase	
Viruses have evolved devices for modulating inflammatory responses	
Proinflammatory cytokines act systemically to promote immunity	
Inflammation is normally downregulated by a series of feedback inhibitory mechanisms	
3-16 Innate Defense Against Viruses: Interferon	86
Interferon provides a critical defense against virus infection	
Interferons are produced by innate immune cells and by infected cells	
3-17 Inhibition of Virus Replication by Interferon	88
Interferon primes cells to allow them to block virus replication	
Viruses can evade the effects of interferon	
Interferons are used to treat some diseases	
3-18 Innate Defense Against Viruses: Induction of Apoptosis	90
Viral replication triggers apoptotic mechanisms in the infected cell	
Death domain receptors can induce death of virus-infected cells	
Viruses have evolved many mechanisms to block apoptosis	

3-19	Intracellular Defenses Targeting Viral Nucleic Acid	92
	Antiviral innate immune defenses include mechanisms directed at viral nucleic acid	
	APOBEC3G is a cytidine deaminase that mutates retroviral genomes	
	RNA interference is an antiviral defense mechanism in plants and probably in animals	
CHAPTER 4 Adaptive Immunity and the Detection of Infection by T Lymphocytes		
4-0	Overview: The Role of T Cells in the Adaptive Immune Response	96
	Adaptive immunity is distinct from innate immunity in three main ways	
	T lymphocytes kill cells infected with viruses and activate B cells and phagocytes	
	The major classes of T lymphocytes are defined by the class of MHC molecule they recognize	
	Dendritic cells provide a bridge between innate and adaptive immunity	
4-1	Dendritic Cells in the Induction of Adaptive Immunity	98
	The function of dendritic cells is to present antigen to naïve T cells	
	Immature dendritic cells comprehensively sample their tissue environment	
	Recognition of infection licenses dendritic cells to activate the effector differentiation of naïve T cells	
4-2	The Structure and Function of MHC Molecules	100
	The major histocompatibility molecules monitor the internal compartments of cells	
	The two classes of MHC molecules have a common architecture but distinct functions in immunity	
4-3	The MHC and Polymorphism of MHC Molecules	102
	The major histocompatibility complex contains genes encoding many molecules with different functions in immunity	
	The classical MHC molecules are highly polymorphic	
4-4	Non-Classical MHC Class I Molecules	104
	The non-classical MHC class I molecules are a diverse group of structural variants on the classical MHC class I molecules	
	The peptide-binding grooves of non-classical MHC class I molecules are adapted to different specialized functions	
4-5	Peptide Binding by MHC Molecules	106
	Peptides bind in distinct ways to MHC class I and MHC class II molecules	
	Variable pockets in the peptide-binding groove anchor specific side chains of the peptide in the groove	
4-6	Antigen Presentation by Classical MHC Class I Molecules	108
	Classical MHC class I molecules display peptides generated in the cytosol	
	Enzyme complexes in the cytosol process proteins into peptides	
	A machinery encoded mainly in the MHC exists in the endoplasmic reticulum to load peptides onto MHC class I molecules	
4-7	Antigen Presentation by MHC Class II Molecules	110
	MHC class II molecules display peptides derived from pathogens internalized by specialized immune cells	
	Antigenic peptides loaded by MHC class II molecules are salvaged from endosomal pathways	
	Peptide loading onto MHC class II molecules is chaperoned	
4-8	Regulation of Antigen Presentation	112
	MHC molecule expression and loading is upregulated by infection	
	MHC class I molecule expression and loading are upregulated by interferons	
	Immune cytokines modulate presentation of peptides by MHC class II molecules	
4-9	Specialized Features of Antigen Presentation by Dendritic Cells	114
	The endosomal pathway is highly regulated in dendritic cells	
	Dendritic cells present peptides derived from exogenous proteins on MHC class I molecules to activate naïve CD8 T cells	

CHAPTER 5 Activation and Effector Actions of T Cells

- 5-0 Overview: T Cell-Mediated Immune Responses** 118
T lymphocytes are central to adaptive immune responses
T cells are activated by strong and sustained binding to their antigen receptors maintained by adhesive interactions
Most T cell responses depend upon the recruitment or activation of other effector cells
T cell actions and numbers are controlled by cytokines
- 5-1 Antigen Recognition by T Cells: Structure of the Receptor** 120
Antigen recognition is mediated by a heterodimeric molecule linked to signaling molecules
There are two kinds of variable receptors for antigen on T cells
The variability of T cell receptors is generated by DNA rearrangements during T cell ontogeny
- 5-2 Antigen Recognition by T Cells: TCR Specificity and Coreceptors** 122
The mature T cell receptor repertoire is determined by the germ-line pool of gene segments and thymic selection
Variable binding orientation and induced fit contribute to the versatility of peptide–MHC binding by $\gamma\delta$ TCR
 $\gamma\delta$ TCR bind invariant ligands
T cells bind to MHC molecules and signal with the help of coreceptors
- 5-3 Signaling from the T Cell Receptor** 124
T cells are sensitive to very small amounts of foreign peptide
TCR signaling is initiated by phosphorylation of receptor ITAM sequences
TCR signaling activates cytoskeletal remodeling and cytokine production
- 5-4 Regulation of TCR Signaling** 126
TCR signaling is dynamically regulated to support T cell activation
The contact between T cells and antigen-presenting cells can result in organized arrays of signaling molecules
- 5-5 T Cell Signaling to the Nucleus and the Role of CD28** 128
TCR signaling leads to activation of proliferation and clonal expansion through induction of cytokine gene transcription
IL-2 gene transcription requires the activation of four transcriptional regulators by the TCR
The costimulatory receptor CD28 amplifies the signal from the TCR
- 5-6 Activation of Naïve T Cells in Secondary Lymphoid Tissue** 130
Activation of naïve T cells occurs when recirculating T cells recognize antigen at the surface of dendritic cells in secondary lymphoid tissues
Naïve T cells are directed into secondary lymphoid tissues by a characteristic combination of homing molecules
T cells that bind strongly to self-peptide–MHC complexes may be inactivated
T cell activation is initiated by lymph-node-resident dendritic cells and stabilized by recruited dendritic cells
- 5-7 Clonal Expansion of Naïve T Cells in Secondary Lymphoid Tissue** 132
Activation of T cells leads to proliferation accompanied by changes in their surface properties
Amplification of T cell numbers on activation is rapid and substantial
The engagement of the TCR and CD28 by peptide–MHC and B7 on the dendritic cell activates a positive feedback loop
- 5-8 Costimulatory Signals in T Cell Proliferation and Survival** 134
Induced modulators of T cell signaling regulate T cell survival and proliferation
Activation of CD8 T cell immunity sometimes requires help from CD4 T cells
- 5-9 Effector T Cell Differentiation** 136
CD4 T cells differentiate into subsets characterized by distinct patterns of cytokine expression
CD8 T cells most commonly differentiate into T_C1 effector cells
T cell differentiation is driven by dendritic cells and cytokines
Most effector T cells disperse to peripheral tissues and accumulate at sites of infection

5-10 The Polarization of T_H1 and T_H2 Cells	138
Polarization of T cell subsets is accompanied by changes in key transcriptional regulators	
The differentiation of T _H 1 cells is mediated by induction of T-bet	
The differentiation of T _H 2 cells is mediated by induction of GATA-3	
5-11 The Functions of T_H1 Cells	140
T _H 1 cells coordinate the destruction and control of pathogens by cytolytic cells and activated phagocytes	
The effects of T _H 1 cells are mediated by cytokines	
T _H 1 cytokines induce B cells to make the types of antibodies that promote destruction of pathogens by phagocytic cells	
T _H 1 cells coordinate a system for antigen elimination but can cause immunopathology	
5-12 The Functions of T_H2 Cells	142
T _H 2 cells coordinate barrier immunity against invading pathogens	
T _H 2 cytokines induce alternatively activated macrophages and activate eosinophils, basophils and mast cells	
T _H 2 cytokines induce B cells to produce antibodies involved in barrier immunity	
Uncontrolled T _H 2 responses can cause immunopathology	
5-13 The Functions of T_{FH} Cells, T_H17 Cells and Induced T_{REG} Cells	144
T _{FH} cells provide help for germinal center B cells	
T _H 17 cells recruit neutrophils to inflammatory sites	
Induced T _{REG} suppress activation of effector T _H cells	
5-14 Effector Functions of CD8 T Cells	146
CD8 T cells are important for the clearance of viruses and bacteria that proliferate in the cytosol	
There are two main mechanisms of CD8 T cell cytotoxicity	
CD8 T cells are protected from their own cytotoxic products	
5-15 T Cell Homeostasis and the Termination of Immune Responses	148
T cell numbers are subject to dynamic control	
Numbers of naïve T cells are maintained by cytokines and stimulation through the TCR	
Effector T cells are programmed to undergo apoptosis	
Inhibitory signals help to limit T cell responses in the periphery	
5-16 T Cell Memory	150
Two types of memory T cells are generated during immune responses	
Memory T cells are maintained by cytokines and stimulation through the TCR, and possibly by antigen	
T cell memory is particularly important for immunity to latent viruses	
Chapter 6 B Cells and Humoral Immunity	
6-0 B Cells and the Importance of Antibodies for Immune Defense	154
Humoral immunity is complementary to cell-mediated immunity	
Antibodies produced by distinct types of B cells are important in early protection from infection and in preventing reinfection	
6-1 Structure of Antibodies	156
Antibodies contain an antigen-binding region and a region that connects to effector mechanisms	
6-2 Structural Properties of Antibodies	158
There are five main kinds of antibodies with distinct functional properties	
Specialized Fc receptors transport antibodies to their sites of action	
6-3 Effector Functions of Antibodies	160
Antibodies can act by neutralization and complement activation	
Fc receptors mediate many of the effector functions of antibodies	
Fc γ R effector function is regulated by the balance of activating and inhibitory receptor action	

6-4 Monoclonal Antibodies	162
Monoclonal antibodies can be produced to capture a single antibody specificity	
Monoclonal antibodies have permitted the identification of most specialized surface molecules of lymphocytes	
Human monoclonal antibodies can be made for therapeutic uses	
6-5 Antibody-Based Methods: Radioimmunoassay and Hemagglutination	164
Antibodies have many uses in diagnostic techniques and research	
Immune complexes of antibodies and complex antigens can generate large insoluble aggregates	
Use of antibodies in the analysis and measurement of biological molecules	
6-6 Antibody-Based Methods: ELISA, Immunoblotting and FACS	166
Antibodies bound to a solid substrate can be detected by the production of colored products	
Antibodies can be used to characterize the properties of subsets of cells	
6-7 Antigen Receptors of B Lymphocytes	168
Membrane forms of immunoglobulin comprise the ligand-binding part of the B cell antigen receptor	
The BCR binds native proteins, glycoproteins and polysaccharides	
The type of Ig heavy chain produced is dependent on the differentiated state of the B cell	
6-8 B Cell Antigen Receptor Signaling and B Cell Coreceptors	170
The BCR signals via ITAM sequences in the Ig α and Ig β subunits	
BCR signaling is modulated by positive and negative coreceptors	
6-9 Early and T Cell-Independent Responses by B Cells	172
Protective antibodies are preexisting and induced	
TI antibody responses are induced by two different mechanisms and are critical to the responses to some pathogens	
6-10 Initiation of T Cell-Dependent Antibody Responses	174
Antigen binding induces follicular B cells to migrate to the boundary between the B cell follicles and the T cell zone	
Antigen bound to the BCR is taken up efficiently, processed and presented to helper T cells	
Helper T cells induce rapid antibody production and the germinal center reaction	
6-11 Hypermutation and Isotype Switching in the Germinal Center	176
The germinal center is a transient structure that supports the production of high-affinity antibodies	
Activation-induced cytidine deaminase mediates class switch recombination and hypermutation of Ig genes	
6-12 Selection and Differentiation of B Cells in the Germinal Center	178
Centrocytes are selected for the ability of their antigen receptors to bind effectively to antigen	
Centrocytes give rise to long-lived plasma cells and memory B cells	
Memory B cells can be rapidly activated to produce higher levels of antibodies	
 CHAPTER 7 Development of Lymphocytes and Selection of the Receptor Repertoire	
7-0 Overview of Lymphocyte Development	182
Lymphocyte development generates cells of many different specificities	
Developing lymphocytes make multiple lineage choices	
Lymphocytes undergo positive and negative selection during their development	
7-1 Antigen Receptor Gene Structure and V(D)J Recombination	184
Antigen receptor loci have multiple gene segments that are rearranged to generate functional genes	
Antigen receptor genes are recombined by a site-specific recombinase that generates antigen receptors of great diversity	

7-2	V(D)J Recombination: Mechanism of Recombination RAG-1 and RAG-2 proteins are lymphocyte-specific components of the V(D)J recombinase Non-homologous end joining recombination components also participate in V(D)J recombination Several mechanisms contribute to the generation of junctional diversity	186
7-3	V(D)J Recombination: Control of which Segments Recombine V(D)J recombination is regulated by modifications to chromatin structure affecting DNA accessibility Often, only one allele of an Ig or TCR locus recombines productively	188
7-4	Phylogenetic Diversity in Antigen Receptor Genes The key elements of the adaptive immune system are found in all jawed vertebrates Sharks have a tandem array of V/D/J/C IgH cassettes Chickens and rabbits make extensive use of gene conversion to generate antibody diversity	190
7-5	B Cell Development in the Bone Marrow Commitment to the B cell lineage involves expression of transcriptional regulators and the RAG-1 and RAG-2 proteins B cell developmental stages are defined primarily by Ig gene rearrangement and Ig chain expression Expression of Ig heavy chain is responsible for driving B cell maturation in the bone marrow Self-reactive immature B cells in the bone marrow arrest maturation and undergo receptor editing	192
7-6	Maturation of B Cells in the Periphery Transitional B cells are deleted or anergized by antigen contact in the periphery Transitional B cells are positively selected into different types of mature B cells BAFF is an important survival factor for transitional and mature B cells	194
7-7	Organization of the Thymus and Early T Cell Development The thymus provides a specialized environment for T cell development Notch is important for commitment to the T cell lineage Stages in T cell development are marked by expression of coreceptors Lineage choice between $\alpha\beta$ T cells and $\gamma\delta$ T cells depends on functional TCR rearrangements Pre-TCR signaling induces commitment to the $\alpha\beta$ TCR lineage	196
7-8	Positive Selection of T Cells in the Thymus $\alpha\beta$ TCR specificity determines cell fate and lineage choice between helper and cytotoxic T cells	198
7-9	Proposed Mechanisms of Positive Selection Double-positive thymocytes may mature by a stochastic/selection or by an instructive mechanism	200
7-10	Negative Selection Thymocytes encountering high-affinity peptide–MHC ligands in the thymus die rapidly Special mechanisms exist to promote self-peptide presentation in the thymus Thymocytes discriminate between positive selection and negative selection based in part on TCR off-rate Some thymocytes encountering high-affinity ligands survive and become regulatory T cells	202
7-11	Diseases Related to Lymphocyte Development Defects or aberrations related to lymphocyte development can lead to immunodeficiency or cancers Genetic defects in lymphocyte development lead to immunodeficiency DNA breaks introduced by V(D)J recombination can lead to lymphomas and leukemias	204

CHAPTER 8 Specialized Lymphocytes in Early Responses and Homeostasis

- 8-0 Overview: Specialized Lymphocyte Populations** 208
Specialized types of lymphocytes can be grouped functionally by their capacity to mediate rapid immune responses
Activation of specialized lymphocytes can be mediated by self or foreign ligands
- 8-1 Natural Killer Cells and their Role in Immunity** 210
Natural killer cells provide early protection from intracellular pathogens, particularly viruses
NK cells are recruited and activated by cytokines released by other cells
NK cells express both activating and inhibitory receptors
NK cells express an array of polygenic and polymorphic receptors specific for host MHC class I
- 8-2 Natural Killer Cell Signaling Pathways** 212
Activating NK cell receptors recognize a variety of ligands
Activating NK cell receptors share conserved signaling pathways
Additional inhibitory signals regulate NK cell activation
- 8-3 NKT Cells** 214
NKT cells are specialized for detection of glycolipid antigens
NKT cells are selected in the thymus and populate the liver, spleen and bone marrow
NKT cells release cytokines rapidly after activation and can mediate defense against bacteria not detected by Toll-like receptors
- 8-4 $\gamma\delta$ T Cells** 216
 $\gamma\delta$ T cells express antigen receptors of limited functional diversity
 $\gamma\delta$ T cells develop early in ontogeny and expand as oligoclonal populations in response to diverse environmental antigens
 $\gamma\delta$ T cells have immediate effector function
 $\gamma\delta$ T cells may have a role in tissue repair later in the immune response
- 8-5 Intraepithelial Lymphocytes and Other Specialized T Cells** 218
Intraepithelial lymphocytes populate the epithelia of bowel and lung
Intraepithelial lymphocytes are cytotoxic
Additional specialized lymphocyte populations exist that recognize non-classical MHC class I molecules that may present microbial components
- 8-6 B1 Cells** 220
B1 cells produce natural antibody, which is present before infection, and T cell-independent antibody that protects body cavities
Natural antibody is synthesized by B1 cells from a preprogrammed repertoire of immunoglobulin genes
B1 cells also participate in first-line IgM and IgA responses to bacteria in the peritoneal and pleural cavities
B1 cells require antigen recognition for their development and are restrained by the inhibitory receptor CD5
- 8-7 Marginal Zone B Cells** 222
Blood is filtered in the spleen via the specialized cells of the marginal zone
Marginal zone B cells make antibody responses to bloodborne infectious agents

CHAPTER 9 The Immune Response to Bacterial Infection

- 9-0 Overview: Bacterial Pathogens and Host Defenses** 226
Commensal bacteria colonize microenvironmental niches and contribute to a healthy epithelial interface
Disrupting the commensal bacterial flora creates an opportunity for pathogenic bacteria
Pathogenic bacteria exploit epithelial dysfunction and subvert innate immunity
Control of pathogenic organisms requires both innate and adaptive immune responses

9-1	Epithelial Barriers and Defenses against Infection	228
	Epithelia present both a physical and an antimicrobial barrier	
	Specialized cell types in epithelia initiate and execute adaptive immune responses	
9-2	Evasion of Epithelial Defenses by Pathogenic Bacteria	230
	Pathogenic bacteria have evolved to traverse epithelial barriers	
	Pathogenic bacteria evade innate defense mechanisms	
	Adaptive immunity is required for protection against pathogenic bacteria	
9-3	Sepsis Syndrome: Bacterial Endotoxin	232
	Sepsis syndrome is a systemic response to invasive pathogens	
	Lipopolysaccharide recognition occurs via the innate immune system	
	Sepsis results from the activation of lipopolysaccharide-responsive cells in the bloodstream	
	Inflammation leads to widespread endothelial cell activation and organ dysfunction	
	Animal models have helped clarify mechanisms of sepsis	
9-4	Sepsis Syndrome: Bacterial Superantigens	234
	Bacterial pyrogenic exotoxins can cause sepsis syndrome	
	Bacterial pyrogenic exotoxins are superantigens and activate T cells expressing unique V β T cell receptors	
	Mouse minor lymphocyte stimulating genes are endogenous retroviral superantigens derived from mouse mammary tumor viruses	
	Mice are susceptible to shock induced by superantigens binding to closely similar V β chains	
9-5	The Immune Response to <i>Streptococcus pneumoniae</i>	236
	<i>Streptococcus pneumoniae</i> is a common cause of bacterial disease, with significant mortality	
	Pneumococci disseminate widely in susceptible hosts	
	Control of pneumococci requires anti-capsular antibody and phagocytes	
9-6	<i>Listeria monocytogenes</i>	238
	<i>Listeria monocytogenes</i> can be lethal to individuals with lowered cell-mediated immunity	
	<i>Listeria</i> is a facultative intracellular organism	
	Host immunity requires a type 1 response with T _H 1 cells and cytotoxic T cells	
	The importance of different components of the immune response to intracellular pathogens can be evaluated in mice infected with <i>Listeria</i>	
9-7	<i>Mycobacterium tuberculosis</i>	240
	Tuberculosis is a major world health problem	
	<i>Mycobacterium tuberculosis</i> establishes chronic infection	
	Control of tuberculosis requires T _H 1-mediated immunity, and disease occurs when immunity is impaired	
	Tuberculosis antigens stimulate $\gamma\delta$ T cells and T cells recognizing CD1	
	Inherited deficiencies in T _H 1 immunity predispose to mycobacteria	
 CHAPTER 10 The Immune Response to Viral Infection		
10-0	Overview: Viral Infectious Strategies	244
	Viruses are ubiquitous and highly diverse	
	Tissue-specific receptors underlie viral tropism	
	Viruses evade immune responses by diverse mechanisms	
10-1	Overview: Innate Strategies Against Viruses	246
	RNA and DNA sensors activate cells to achieve an antiviral state and to recruit host immune cells	
	Plasmacytoid dendritic cells are specialized producers of IFN- α	
	NK cells contribute to early antiviral immunity	
10-2	Overview: Adaptive Immune Strategies Against Viruses	248
	Viral infections can activate massive oligoclonal expansion of CD8 T cells	
	Antibodies are critical for the control of many viruses	

10-3 Subversion of Immune Mechanisms by Viruses	250
Pathogenic viruses have captured genes that target cellular apoptosis, interferon and innate inflammation, and the display of peptide-MHC class I complexes	
10-4 Origin and Structure of Human Immunodeficiency Virus	252
AIDS originated as a zoonotic disease introduced into humans from primates	
HIV is a retrovirus in the lentivirus subfamily	
10-5 Tropism of Human Immunodeficiency Virus	254
HIV crosses mucosal barriers, establishes early viremia and targets CD4 T cells	
Activated CD4 T cells drive viral replication while HIV subverts immune detection	
10-6 AIDS Progression and Immune Control of HIV	256
Progression to AIDS is determined by the viral set-point	
Control of HIV is mediated by CD4 T cell-dependent cytotoxic CD8 T cells	
10-7 Influenza Virus: Antigenic Shift and Drift	258
Influenza viruses cause recurrent epidemics and occasional pandemics	
Influenza A viruses are avian viruses that adapt to humans	
Antigenic drift and shift in HA and NA underlie influenza A epidemics and pandemics	
10-8 Influenza Virus: Innate and Adaptive Immunity	260
Innate immunity to influenza depends on type 1 interferons	
Influenza virus infection is cleared by cytotoxic CD8 T cells	
Viral protection is mediated by neutralizing antibody	
10-9 Lymphocytic Choriomeningitis Virus (LCMV)	262
Lymphocytic choriomeningitis virus (LCMV) is extensively used in mouse models of immunity to viruses	
LCMV is a noncytopathic virus that establishes persistent infection in mice	
Acute infection, immune exhaustion and immune tolerance can be studied using LCMV in inbred mice	

CHAPTER 11 The Immune Response to Fungal and Parasitic Infection

11-0 Overview: The Scope of Fungal and Parasitic Infections	266
Pathogenic fungi can be commensals, ubiquitous environmental molds, or geographically endemic species	
Phagocytes and dendritic cells express receptors for fungal constituents	
Host defense against fungi relies on epithelial barriers and phagocyte activation facilitated by T _H 1 and T _H 17 cells	
Protozoa are a diverse group of systemic and intestinal pathogens	
Protozoan pathogens induce disease-protective, but not sterile, immunity	
Helminths constitute a diverse group of systemic and intestinal pathogens	
Helminths cause chronic infections and incomplete immunity	
11-1 <i>Candida albicans</i>	268
<i>Candida albicans</i> is a commensal and opportunistic pathogen of humans	
<i>Candida albicans</i> virulence is linked to the ability to grow vegetatively in morphologically distinct forms	
Phagocyte oxidative killing is important in <i>Candida</i> host defense	
Mouse models exist for both systemic and mucosal <i>Candida</i> infection	
11-2 <i>Pneumocystis</i>	270
<i>Pneumocystis</i> is an important cause of pneumonia in immunocompromised hosts	
<i>Pneumocystis</i> comprises diverse host-specific strains of unusual fungi	
<i>Pneumocystis</i> infection of mice comprises an experimental model for studying the immune response	
11-3 <i>Leishmania major</i>	272
<i>Leishmania major</i> is a protozoan pathogen transmitted by sandflies	

Leishmania parasites invade macrophages without inducing inflammatory cytokines and reside in a phagolysosomal compartment

Immunity to *Leishmania* requires T_H1 cells, and persistence is maintained by regulatory T cells

Some strains of mice develop an aberrant and fatal T_H2 response to *L. major*

11-4 *Nippostrongylus brasiliensis* 274

Nippostrongylus brasiliensis is a roundworm that matures during migration through the lung to the small intestines

Nippostrongylus induces a protective T_H2-associated immune response

Recruitment of immune cells to tissue and activation of worm expulsion reflect compartmentalized signals via the IL-4R α chain and STAT6

11-5 Schistosomiasis 276

Schistosomes are a widespread human trematode infection of the abdominal vasculature

Host immune responses are modulated over the course of infection and culminate in a highly polarized type 2 granulomatous response to eggs

Type 2 granulomatous responses benefit the host and the parasite until fibrosis compromises the function of critical organs of the body

Mouse models for *S. mansoni* infection and for type 2 granulomatous responses reveal distinct roles for T_H2 cytokines

CHAPTER 12 Tolerance and Autoimmunity

12-0 Overview: Tolerance and Autoimmunity 280

Immune tolerance is established and maintained by both central and peripheral mechanisms

Failures in self tolerance lead to autoimmune disease

Some autoimmune diseases are treated by blocking molecules of the immune system

12-1 Central Tolerance 282

Highly self-reactive B cells and T cells are eliminated during their development

The thymus has specialized mechanisms for expressing self antigens

12-2 Peripheral Tolerance Mechanisms 284

Central tolerance is insufficient to generate fully effective immune tolerance

B cells are anergized by antigen contact in the absence of T cell help

Dendritic cells constitutively present self antigen without costimulation or cytokines and induce abortive activation of self-reactive T cells

12-3 Regulatory T Cells 286

Regulatory T cells are a distinct type of effector T cells that suppress T cell immune responses

Regulatory T cells are essential for avoiding pathological immune responses

12-4 Autoimmune Diseases: General Principles 288

Many diseases result from immune attack on uninfected tissues

Autoimmunity is typically the result of genetic susceptibility and an environmental trigger

Molecular mimicry and epitope spreading may be important for autoimmunity

12-5 Organ-Specific Autoimmunity: Antibody-Mediated Diseases 290

Antibodies against self components can cause disease

Autoantibodies can recognize cells or extracellular matrix and induce pathology

Autoantibodies can cause disease by affecting the function of the target autoantigen

12-6 Organ-Specific Autoimmunity: Cell-Mediated Diseases 292

Cell-mediated autoimmune diseases attack a variety of organs

T cell costimulation is likely to be important for progression of cell-mediated autoimmune diseases

12-7 Organ-Specific Autoimmunity: Animal Models 294

Animal models of cell-mediated autoimmune disease have been informative about the mechanisms of autoimmunity

NOD mice spontaneously develop type 1 diabetes

Experimental allergic encephalitis has many similarities to multiple sclerosis	
Molecular mimicry is responsible for a virus-induced autoimmune eye disease	
12-8 Systemic Autoimmunity: Lupus	296
Systemic autoimmune diseases include SLE and rheumatoid arthritis	
SLE is characterized by the production of high levels of antibodies against nuclear components	
Mouse models provide insight into the genetic susceptibility of SLE	
12-9 Systemic Autoimmunity: Rheumatoid Arthritis	298
Rheumatoid arthritis is a common systemic autoimmune disease	
CHAPTER 13 Allergy and Hypersensitivity	
13-0 Overview: Causes and Nature of Hypersensitivity Reactions	302
Hypersensitivity diseases result from excessive responses to non-infectious environmental antigens	
Hypersensitive responses occur only after prior sensitization	
Antigens that elicit hypersensitive responses can be haptens that modify host proteins	
The time course of a hypersensitive reaction depends on the mechanism	
13-1 Mast Cells and Allergic Reactions	304
Immediate-type hypersensitivity reactions are initiated by binding of allergen to IgE on sensitized mast cells	
Cytokines and lipid mediators are responsible for prolonged effects of mast cell triggering	
Mast cells induce an inflammation rich in basophils, eosinophils and T _H 2 cells	
13-2 Allergic Diseases	306
Allergic diseases have both environmental and genetic causes	
Manifestations of allergic diseases are site-dependent	
Anaphylaxis is a systemic response to an allergen	
Allergic rhinitis occurs in response to allergens in the nasal passages	
Food allergy is commonest in young children	
Allergies often manifest in skin	
13-3 Asthma	308
Asthma is a respiratory disease of increasing prevalence	
Asthma reflects dysregulated T _H 2-associated immunity	
Asthma represents an interaction of allergens and genetics	
Mouse models mimic aspects of acute airway inflammation in human asthma	
13-4 IgG-Mediated Immune Pathology	310
A variety of hypersensitivities and autoimmune diseases exhibit immune pathology caused by IgG	
Hemolytic anemias can be caused by Ig produced in response to infection or materno-fetal Rhesus incompatibility	
Immune complexes can cause damage to joints, blood vessels and kidneys	
13-5 Delayed-Type Hypersensitivity Reactions	312
Delayed-type hypersensitivity reactions reflect the actions of T _H 1 and cytotoxic cells	
Contact sensitivity to urushiol is an inflammatory response to haptened proteins	
Celiac disease is a delayed-type hypersensitive response to gliadin peptides from dietary wheat	
Delayed-type hypersensitive reactions are used to test for T cell function or for the presence of specific infections or hypersensitivities	
13-6 Inflammatory Bowel Disease	314
Crohn's disease and ulcerative colitis are chronic intestinal inflammatory diseases	
Crohn's disease represents a dysregulated mucosal immune response	
Ulcerative colitis represents a dysregulated T _H 2-associated mucosal immune response	
Animal models for inflammatory bowel disease are diverse	

CHAPTER 14 Transplantation Immunology, Tumor Immunity and Vaccination

14-0 Overview: Boosting and Suppressing Immune Responses	318
Modulating the immune response is often important for improving human health	
Vaccination is an important tool in the fight against infectious diseases	
Immunosuppression is often required for the success of cell and organ therapies	
14-1 Immunodeficiencies	320
Genetic deficiencies in components of immunity often increase susceptibility to particular infections	
Some immunodeficiencies are acquired	
14-2 Features of Successful Vaccines	322
Successful vaccines depend on production of antibody	
Adjuvants are required to potentiate and amplify immune responses to vaccines	
14-3 Engineering Vaccines for Safety and Efficacy	324
Declining incidence of infectious disease may encourage a decline in vaccination	
The combined diphtheria/tetanus/pertussis vaccine is based on toxoids with added adjuvants	
Conjugate vaccines activate T cell help for B cell responses to polysaccharides	
The route of immunization is important in determining the type of protection elicited	
14-4 Vaccines for Chronic Infections and Cancer	326
Persistent pathogens present a special challenge for vaccine design	
HIV escapes both neutralizing antibodies and cytotoxic T cells	
Malaria is a complex target for recombinant vaccines	
Cancer can be prevented by vaccines against tumorigenic pathogens	
14-5 Tumor Immunity	328
The immune system can eliminate many tumors before they become evident	
Both innate and adaptive immune cells can detect cancer cells	
Normal, mutant and foreign proteins can serve as tumor antigens	
Cytokines, antibodies and T cells are all potentially useful for cancer immunotherapy	
14-6 Transplants and Transplantation Antigens	330
Immunological rejection is a major barrier to transplantation	
Transplantation antigens include the blood group antigens, MHC and minor histocompatibility antigens	
14-7 Transplant Recognition and Rejection	332
Organ transplant rejection occurs by three different mechanisms	
T cells can be stimulated by dendritic cells of the graft or of the host	
14-8 Immunosuppression of Transplant Recipients	334
Transplantation is successful because of immunosuppressive drugs and antibodies	
Steroids, antiproliferative agents and calcineurin inhibitors are the mainstays of immunosuppression for transplantation	
Anti-CD3 antibodies are used to combat rejection episodes	
Glossary	337
References	355
Index	370